

In vitro antioxidant properties of *Portulaca quadrifida* Linn.

¹ Manivannan E*, ² Kothai R and ¹ Arul B.

¹ Department of Pharmacology, Vinayaka Mission's Kirupanandha Variyar Medical College, Vinayaka Missions University, Seerapadi Salem, Tamilnadu, India.

² Department of Pharmacology, Vinayaka Mission's College of Pharmacy, Salem.

*Corresponding Author: E-Mail: manipoo73@gmail.com

Received: 25 June 2015, Revised and Accepted: 06 July 2015

ABSTRACT

The purpose of this work is to assess the *in vitro* antioxidant activity of ethanolic extract of *Portulaca quadrifida* Linn. by using the following analytical methods: 1, 1-diphenyl-2-picrylhydrazyl (DPPH•) free radical scavenging, nitric oxide radical inhibition, superoxide anion (O₂^{•-}) radical scavenging, inhibition of xanthine oxidase activity and the inhibition of oxidation of β-carotene. Results of ethanolic extract of *Portulaca quadrifida* were compared with the activity of respective standard drugs and the extract was found to be equipotent with the standards in all the tested methods. The flavonoids present in the *Portulaca quadrifida* may be believed to be responsible for its antioxidant activity.

Keywords: *Portulaca quadrifida*; Antioxidant; β-carotene oxidation; DPPH; Nitric oxide; superoxide anion; xanthine oxidase.

1. INTRODUCTION

Oxidative damage in the human body plays an important causative role in disease initiation and progression [1]. Damage from reactive oxygen species (ROS) including free radicals has been linked to some neurodegenerative disorders (alzheimer's disease and parkinson's) [2] and cancers [3]. ROS include free radicals such as superoxide anion radicals (O₂^{•-}), hydroxyl radicals (OH•) and non-free-radical species such as H₂O₂ and singlet oxygen (¹O₂). These molecules are exacerbating factors in cellular injury and aging process [4, 5]. Recently, ROS have also been shown to play a critical role in the development of acute experimental gastric lesions induced by stress, ethanol and NSAIDs [6-8]. Exogenous sources of ROS include tobacco smoke, certain pollutants, organic solvents and pesticides [9]. ROS also induce some oxidative damage to biomolecules such as lipids, nucleic acids, proteins and carbohydrates. Damage of these biomolecules causes aging, cancer, and many other diseases [9]. As a result, ROS have been implicated in more than one hundred diseases, including malaria, heart disease, stroke, arteriosclerosis, diabetes, cancer and gastric ulcers [10-12].

The human body has inherent mechanisms to reduce free radical injury by

endogenous enzymes such as superoxide dismutase, glutathione peroxidase, catalase. Exogenous free radical scavengers include vitamin E and vitamin C [13, 14]. These vitamins protect the human body and retard the progress of many chronic diseases as well as lipid oxidative rancidity in foods [15-17]. Antioxidant defense systems, including antioxidant enzymes which are found in foods and drugs are important in the prevention of the toxic ROS effects [14, 18, 19]. Hence, the search for exogenous antioxidants is continued.

The most commonly used antioxidants at the present time are butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), propyl gallate and *tert*-butylhydroquinone [20]. However, BHA and BHT have been suspected of being responsible for liver damage and carcinogenesis [20, 21]. Therefore, the development and utilization of more effective antioxidants is desired [5, 22]. Much attention has been focused on the antioxidative compounds present in plants because of safety concerns associated with synthetic antioxidants [23].

Portulaca quadrifida Linn. (portulacaceae) a prostrate fleshy annual or stoloniferous perennial herb with somewhat base but sometimes with simple main stems, 5-40 cm

tall and generally widespread in warm countries. It is a small diffused, succulent, annual herb found throughout the tropical parts of India. It is used as a vegetable and also used for various curative purposes. It is said to be useful in asthma, cough, urinary discharges, inflammations and ulcers. A poultice of the plant is applied in abdominal complaints, erysipelas and haemorrhoids [24]. *Portulaca quadrifida* Linn. has been reported to possess antifungal activity against *Aspergillus fumigates* and *Candida albicans* [25] and the neuropharmacological activities were reported by Syed *et al* [26]. A review of literature afforded no information on the antioxidant activity of this plant. So the present study is therefore an attempt to assess the efficacy of this indigenous herb for its *in-vitro* antioxidant activity by various methods.

2. EXPERIMENTAL

2.1. Materials

2.1.1. Plant material

The aerial parts of the plant *Portulaca quadrifida* were collected from the foothill of Yercaud, Salem, in the month of June 2013 and cleaned to remove the debris. The collected plant was identified and authenticated by a botanist Dr. A. Marimuthu, Department of Botany, Government Arts College, Attur. A voucher specimen (PQM-1) has been kept in our museum for future reference. The plant parts were dried at room temperature for 10 d and coarsely powdered with the help of a hand-grinding mill and the powder was passed through sieve No. 60.

2.1.2. Chemicals

1, 1-Diphenyl-2-picryl hydrazyl (DPPH) was obtained from Sigma-Aldrich Co., St. Louis, USA. Rutin was obtained from Acros Organics, NJ, USA. Naphthyl ethylene diamine dihydrochloride (NEDD) was from Roch-Light Ltd., Suffolk, UK. Allopurinol was from GlaxoSmithKline, Philadelphia, USA. Superoxide dismutase (SOD) was from Shenzhen Jenlong Biotechnology Co., Ltd, China. Butylated hydroxy anisole (BHA) was from SD Fine Chemicals Ltd., Mumbai, India. Sodium nitroprusside was from Ranbaxy Laboratories Ltd., Mohali, India. All chemicals used were of analytical grade.

2.2. Methods

2.2.1. Preparation of the extract

The powder of aerial parts of *P. quadrifida* was extracted separately by continuous hot extraction process using soxhlet apparatus with different solvents in increasing order of polarity from petroleum ether, chloroform, acetone, alcohol, to finally chloroform:water. After

extraction, the extracts were concentrated under reduced pressure in tared vessel. The marc of crude drug powder was then once again subjected to successive extraction with other solvents and the extractive values were calculated with reference to the air-dried drug. The dry extracts were subjected to various chemical tests to detect the presence of different phytoconstituents.

2.2.2. 1,1-Diphenyl-2-picryl-hydrazil (DPPH) freeradical-scavenging activity

The stable free radical scavenging effect of ethanolic extract of *P. quadrifida* (EPQ) was assessed by the decoloration of a methanolic solution of DPPH according to the Blois' method [27]. Test samples (20 -100 µg/ml) were dissolved in 0.1 ml DMSO and then added to 0.1 ml of 0.1 mM DPPH in methanol. The mixture was shaken vigorously and allowed to stand for 10 minutes at room temperature (30°C) in the dark. The absorbance at 517 nm by DPPH was measured spectrophotometrically (SPECTRA_{MAX} PLUS³⁸⁴, Molecular Devices, USA) at 517 nm. Ascorbic acid was used as a positive control. The degree of discoloration indicates the scavenging efficacy of the extract. The capability to scavenge the DPPH radical was calculated using the following equation:

$$\text{DPPH}^{\bullet} \text{ scavenging effect (\%)} = \frac{(A_0 - A_1 / A_0) \times 100}{100}$$

Where in A_0 is the absorbance of the control reaction and A_1 is the absorbance in the presence of EPQ [22, 28].

2.2.3. Nitric oxide radical inhibition assay

Sodium nitroprusside in aqueous solution at physiological pH spontaneously generates nitric oxide, which interacts with oxygen to produce nitrite ions. These ions can be estimated by the use of Griess Illosvoy reaction [29]. In the present investigation, Griess Illosvoy reagent is modified by using naphthyl ethylene diamine dihydrochloride (0.1 %, w/v) instead of 1-naphthylamine (5 %). Nitric oxide scavengers compete with oxygen leading to reduce the production of nitric oxide [30]. The reaction mixture (3 ml) containing sodium nitroprusside (10 mM, 2 ml), phosphate buffer saline (0.5 ml) and extract (EPQ) or standard (Rutin) solution (0.5 ml) was incubated at 25° C for 150 minutes. After incubation, 0.5 ml of the reaction mixture containing nitrite was pipetted and mixed with 1 ml of sulfanilic acid reagent (0.33 % in 20 % glacial acetic acid) and allowed to stand for 5 minutes for completing diazotization. Then, 1 ml of naphthyl ethylene diamine dihydrochloride was added, mixed, and allowed to stand for 30 minutes at 25°C. A pink colored chromophore was formed

in diffused light. The absorbance of these solutions were measured spectrophotometrically (SPECTRA_{MAX} PLUS³⁸⁴, Molecular Devices, USA) at 540 nm against the corresponding blank solutions. IC₅₀ value is the concentration of sample required to inhibit 50 % of the nitric oxide radical.

2.2.4. Scavenger effect on superoxide anion

Superoxide anion was generated in vitro using the method described by Paoletti et al [31]. The assay mixture contained in a total volume of 1 ml, 100 mM triethanolamine-diethanolamine buffer, pH 7.4, 3 mM NADH, 25 mM/12.5 mM EDTA/MnCl₂, 10 mM β-mercapto-ethanol and EPQ at different concentrations (0.15, 0.3, 0.6, 1.2, 2.4 μg/ml). After a 20 min incubation at 25° C, the decrease in absorbance was measured spectrophotometrically (SPECTRA_{MAX} PLUS³⁸⁴, Molecular Devices, USA) at λ = 340 nm. SOD (80 mU/ml) was used as a standard to compare the activity of EPQ.

2.2.5. Xanthine oxidase activity inhibition

Xanthine oxidase activity was evaluated spectrophotometrically as previously reported by following the formation of uric acid at λ = 292 nm (εM = 9.2×10³) [32]. The assay mixture contained, in a final volume of 1 ml, 50 mM phosphate buffer pH 7.8, 25 μM solution of xanthine and 24 mU xanthine oxidase (specific activity 1 U/mg of protein). Different concentrations of EPQ (25, 50, 100, 200 μg/ml) were added to samples before the enzyme and their effect on the generation of uric acid was used to calculate regression lines. The results were expressed as a percentage of inhibited enzyme activity with allopurinol (30 μM) was used as a standard.

2.2.6. β-Carotene oxidation

The assay system used to evaluate the potential of EPQ to suppress β-carotene oxidation was the coupled oxidation of β-carotene and linoleic acid described by Vaya et al [33] with two minor modifications. The plant powder was dissolved in water instead of alcohol to a final concentration of 100 μg/mL and spectrophotometric readings were taken (SPECTRA_{MAX} PLUS³⁸⁴, Molecular Devices, USA) at 454 nm. The assay was performed at 50° C. The readings of the rate of β-carotene bleaching were recorded at 20 minutes intervals for 2 hours against water blank. BHA (100 μg/mL) was served as a positive control. This method allows evaluation of the antioxidative activity of a preparation over a time scale. This assay was repeated three times.

3. RESULTS

Since the DPPH test can accommodate a large number of samples in a short period of time and it is sensitive enough to detect natural compounds at low concentrations, it was used in the present study for a primary screening of EPQ for its free radical-scavenging activity. This assay provides information on the reactivity of test compounds with a stable free radical. Because of its odd electron, DPPH gives a strong absorption band at 517 nm in visible spectroscopy (deep violet color). As this electron becomes paired off in the presence of a free radical scavenger, the absorption vanishes, resulting in stoichiometric decolorization with respect to the number of electrons taken up [27].

The EPQ exhibited DPPH free radical scavenging activity in a concentration-dependent manner (Figure 1). Increasing concentrations of EPQ were taken and the following percent inhibition was observed subsequently at 20 μg/ml (65.17 ± 1.87), 40 μg/ml (70.10 ± 2.23), 60 μg/ml (76.24 ± 1.12), 80 μg/ml (85.78 ± 2.78) and 100 μg/ml (93.96 ± 2.91) with Ascorbic acid equaling 100 μM (94.98 ± 1.11). Significant DPPH radical scavenging activity was evident at all tested concentrations of the extract.




Figure - 1: Scavenger effect of EPQ, at different concentrations (20, 40, 60, 80 and 100 μg/ml), on the stable 1,1-diphenyl-2-picrylhydrazyl radical (DPPH). (Results are expressed as percentage decrement of absorbance at 517 nm with respect to control. Ascorbic acid (100μM) was used as a standard. Each value represents the mean±S.D. of four experiments, performed in duplicate).

Table - 1: In vitro antioxidant activity of P. quadrifida against Nitric oxide method

Test compounds	IC ₅₀ values ± S.E.* (μg/ml)
EPQ	424.25±7.97
Rutin	467.92±9.85

*. Average of 10 determinations

Among the extract and standard tested for antioxidant activity using the nitric oxide radical inhibition method, the EPQ showed better activity. It showed the IC₅₀ value of 424.25±7.97 µg/ml, where as the standard, Rutin showed 467.92±9.85 µg/ml (Table 1).

As DPPH is a synthetic radical, we therefore also investigated the superoxide anion scavenging capacity of this extract using the method of Paoletti et al, which excludes the Fenton-type reaction and the xanthine/xanthine oxidase system [31]. In this assay, EPQ showed a dose-dependent superoxide scavenging effect (Figure 2).




Figure - 2: Scavenger effect of EPQ, at different concentrations (0.15, 0.3, 0.6, 1.2, 2.4 µg/ml), on O₂⁻ expressed as percentage of inhibition of NADH oxidation. (Superoxide dismutase (SOD) (80 mU/ml) was used as a standard. Each value represents the mean±S.D. of four experiments, performed in duplicate).




Figure - 3: Effect of ethanolic extract from EPQ, at different concentrations (25, 50, 100, 200 µg/ml), on xanthine oxidase activity. (Results are expressed as percentage of inhibition of xanthine oxidase activity with respect to control. Allopurinol (30 µM) was used as a standard. Each value represents the mean±S.D. of four experiments, performed in duplicate).

The effect of EPQ on xanthine oxidase activity is shown in figure 3. Results are expressed as a percentage of inhibition of xanthine oxidase

activity with respect to control. The extract showed a dose-dependent inhibition of uric acid formation. Under our experimental conditions, 200 µg/ml of extract was corresponded with 30 µM of allopurinol activity.

The antioxidant potential of the extract was evaluated by determining the ability of 100 µg/mL extract to inhibit β-carotene oxidation. The rate of β-carotene oxidation was inhibited up to 84.95 %, whereas BHA almost completely inhibited the rate of oxidation (99.02 %). Typical curves showing the rate of β-carotene oxidation (bleaching) are presented in figure 4.




Figure - 4: Effect of 100 µg/mL *P. quadrifida* extract on the oxidation of β-carotene. (Data are shown as mean±S.D).

4. DISCUSSION

Oxidants are involved in many human diseases and the aging processes. Chronic damage associated with the development of aging can generate destructive oxidants and oxygen free radicals, which are very toxic to tissues and may result in further tissue necrosis and the cellular damage. Cellular mechanisms and external factors involved in the production of oxidative stress include the inflammatory response, free radical leak from mitochondria, auto-oxidation of catecholamines, xanthine oxidase activation, prooxidant activities of toxins such as CCl₄ and exposure to ionizing radiation [34]. Aerobic cells are endowed with extensive antioxidant defense mechanisms including both low molecular weight scavengers, such as α-tocopherol, cysteine, β-carotene, reduced glutathione, ascorbic acid and enzymatic systems, such as superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GSH-Px), glutathione reductase (GSH-Red) and glucose-6-phosphate dehydrogenase (G6PD), which counteract the damaging effects of reactive oxygen species [35]. However, when the balance between these reactive species and antioxidants are altered, a state of oxidative stress results, possibly leading to permanent cellular damage. Our results demonstrate that EPQ exhibited an interesting antioxidant activity in cell-free systems. It was able to quench the synthetic DPPH

radical and exhibited a SOD-like effect, inhibiting $O_2^{\cdot-}$ formation in a dose-dependent manner [34].

It is reported that green tea tannins are able to protect renal cells against ischemia reperfusion injury [36], characterized by an overproduction of $O_2^{\cdot-}$ due to both an electron leak in the mitochondrial respiratory chain and the conversion of xanthine dehydrogenase to xanthine oxidase [37], which produces $O_2^{\cdot-}$ when it oxidizes xanthine into uric acid. This study, therefore, also considered a possible inhibitory action of the extract on the primary function of this enzyme. The results obtained showed that EPQ may determine a dose-dependent inhibition of xanthine oxidase activity.

The IC_{50} values observed were found to be slightly lower than that of standard rutin in nitric oxide radical inhibition assay. The results clearly indicated that the extract was found to be more effective in scavenging the DPPH free radical when compared to the nitric oxide radical. We also observed the extract inhibited the β -carotene oxidation at 100 μ g/mL concentration.

The first step in investigating the antioxidant potential of EPQ involved defining the relationship between concentration and effect by evaluating its ability to inhibit oxidative processes. Initially, we used a single concentration of extract against all the methods. Having demonstrated that the extract was effective at inhibiting this oxidative process, we then proceeded to confirm this finding with various concentrations of the extract that inhibited the oxidation.

On the basis of these experiments, it could be concluded that the ethanolic extract of *Portulaca quadrifida* Linn. can act as primary and secondary antioxidant. It scavenges free radicals and therefore inhibits the lipid peroxidation and may have beneficial effect on prevention of diseases, where reactive oxygen species are involved. Antioxidant properties of its flavonoid compounds can be at the origin of these effects, but further in vivo experiments are planned to verify the relationship between chemical composition and antioxidant activity.

5. REFERENCES

- Jacob RA and Burri BJ. **American Journal of Clinical Nutrition**, 1996; 63: 985-990.
- Youdim K and Joseph JA. **Free Radical Biology and Medicine**, 2001; 30: 583-594.
- Goodwin JS, Brodwick M. *Clinics in Geriatric Medicine* 1995; 11: 577-589.
- Halliwell B and Gutteridge JM. **Free Radicals in Biology and Medicine**, Clarendon Press, Oxford, 1989: 23-30.
- Gu'lcin 'I, Bu'yu'kokurog'lu MEM, Oktay M, Ku'freviog'lu and O' 'I. **Journal of Pineal Research**, 2002; 33: 167-171.
- Banarjee RK. **Biochimica et Biophysica Acta**, 1990; 1034: 275-279.
- Das D, Bandyopadhyay D, Bhattacharjee M and Banerjee RK. **Free Radical Biology and Medicine**, 1997; 23: 8-18.
- Miura T, Muraoka S and Fujimoto Y. **Biochemical Pharmacology**, 2002; 63: 2069-2074.
- Gulcin I, Berashvili D and Gepdiremen A. **Journal of Ethnopharmacology**, 2005; 101: 287-293
- Halliwell B. **American Journal of Medicine**, 1991; 91: 14-22.
- Hertog MGL, Feskens EJM, Hollman PCH, Katan MB and Kromhout D. **Lancet**, 1993; 342: 1007-10014.
- B'uy'ukokuro'glu ME, Gu'lcin 'I, Oktay M, Ku'freviog'lu, O' 'I. **Pharmacological Research**, 2001; 44: 491-495.
- Yoshikawa T, Minamiyama Y, Ichikawa H, Takahashi S, Naito Y and Kondo M. **Free Radical Biology and Medicine**, 1997; 23: 243-250.
- Mates JM, Perez-Gomez C and Nudez de Castro I. **Clinical Biochemistry** 1999; 32: 595-603.
- Kinsella JE, Frankel E, German B and Kanner JI. **Food Technology**, 1993; 47: 85-89.
- Lai LS, Chou ST and Chao WW. **Journal of Agricultural and Food Chemistry**, 2001; 49: 963-968.
- Gu'lcin 'I, Bu'yu'kokurog'lu ME, Ku'freviog'lu and O' 'I. **Journal of Pineal Research**, 2003; 34: 278-281.
- Ames BN, Shigenaga MK and Hagen TM. **Oxidants, antioxidants, and the degenerative diseases of aging. In: Proceedings of the National Academy of Sciences of the United States of America**, 1993; 90: 7915-7922.
- Bafna PA and Balaraman R. **Journal of Ethnopharmacology**, 2004; 90: 123-127.
- Sherwin ER. In: Branen AL, Davidson PM, Salminen S. (Eds.), **Food Additives**. Marcel Dekker Inc, New York, 1990: 139-193.
- Grice HC. **Food and Chemical Toxicology**, 1986; 24: 1127-1130.
- Oktay M, Gu'lcin 'I, Ku'freviog'lu and O' 'I. **Lebensmittel Wissenschaft and Technologie**, 2003; 36: 263-271

23. Imaida K, Fukushima S, Shirai T, Ohtai M, Nakanishi K and Ito N. **Carcinogenesis**, 1983; 4: 7895-99.
24. Kirtikar and Basu. **Indian Medicinal Plants**, 2001; 2: 333-335.
25. Hoffman BR, Delas Alas, Blanco K, Wiederhold N, Lewis RE and Williams L. Scening of Antibacterial and Antifungal Activities of Ten Medicinal Plants from Ghana. **Pharm. Biol.**, 2004; 42(1): 13-17
26. Syed Kamil M, LiyakhaT Ahmed MD and Paramjyothi S. Neuropharmacological Effects of Ethanolic Extract of *Portulaca quadrifida* Linn. In Mice. **International Journal of PharmTech Research**, 2010; 2(2): 1386-1390.
27. Blois MS. **Nature**, 1958; 26: 1199–1200.
28. Gu'lcin 'I, Bu'yu'kokurog'lu ME, Oktay M, Ku'freviog'lu and O' 'I. **Journal of Ethnopharmacology**, 2003; 86: 51–58.
29. Garrat DC. **The Quantitative Analysis of Drugs**. 3rd ed., Chapman and Hall Ltd., Tokyo, Japan, 1964; 456–458.
30. Marcocci L, Packer L, Sckaki A and Albert GM. **Methods Enzymology**, 1994; 234: 462–475.
31. Paoletti F, Aldinucci D, Mocalli A and Caparrini A. **Analytical Biochemistry**, 1986; 154: 536–541.
32. Russo A, Acquaviva R, Campisi A, Sorrenti V, Di Giacomo C, Virgata G and Vanella A. **Cell Biology and Toxicology**, 2000; 16: 91–98
33. Vaya I, Belinky PA and Aviram M. **Free Radical Biology and Medicine**, 1997; 23: 302–313.
34. Russo A, Cardile V, Lombardo L, Vanella L, Vanella A and Garbarino JA. **Journal of Ethnopharmacology**, 2005; 100: 323–332
35. Halliwell B. **Lancet**, 1994; 344: 721–724.
36. Yokozawa T, Dong E, Chung HY, Oura H and Nakagawa H. **Bioscience Biotechnology and Biochemistry**, 1997; 61: 204–206.
37. Wernes SW and Lucchesi R. **Trends in Pharmacological Sciences**, 1990; 11: 161–166.