

1-(3, 4-dichlorobenzenesulfonyl)-3-methyl-1*H*-imidazolium chloride: An efficient catalyst for the synthesis of 1, 8-dioxo-octahydroxanthenes under microwave irradiation

Kesturu Venkatarayappa Jagannath and Krishnamurthy Gantigaiah*.

Department of Studies in Chemistry, Central College Campus, Bangalore University, Bengaluru, Karnataka, India.

*Corresponding Author: E-Mail: krismurg@gmail.com

Abstract

The reaction of 5,5-dimethyl-1,3-cyclohexanedione with various aromatic aldehydes to give corresponding 1,8-dioxo-octahydroxanthenes in aqueous medium has been reported. These reactions were carried out under microwave irradiation and catalyzed by 1-(3, 4-dichlorobenzenesulfonyl)-3-methyl-1*H*-imidazolium chloride. A plausible mechanism for the synthesis of 1,8-dioxo-octahydroxanthenes catalyzed by [3, 4-dcbsmim][Cl] in water has been given.

Keywords: Aldehydes, 1-(3, 4-dichlorobenzenesulfonyl)-3-methyl-1*H*-imidazolium chloride ([3,4-dcbsmim][Cl]), 5,5-dimethyl-1,3-cyclohexanedione, 1,8-dioxo-octahydroxanthenes, Microwaves, Water.

1. INTRODUCTION

Xanthenes constitute an important class of biologically active compounds because of its wide range of biological and pharmaceutical properties such as anti-inflammatory, antidepressants and antimalarial agents [1]. They act as sensitizers in photodynamic therapy (PDT) for destroying the tumor cells [2]. Xanthenes are also used in the dye industry [3], in cosmetics and pigments [4], in laser technology because of their spectroscopic properties [5], etc. Xanthenes based dyes are extracted naturally from soil and plants such as *Indigofera longiracemosa* [6]. They are used as pH sensitive fluorescent materials for the visualization of biomolecular assemblies [7] and also as antagonists for the paralyzing action of zoxazolamine [8].

Due to their biological and pharmacological importance, several methods have been reported for the synthesis of Xanthenes employing Lewis acids as catalysts to accelerate the condensation reaction of aldehydes with 5,5-dimethyl-1,3-cyclohexanedione in the presence of [1-(chloromethyl)-4-fluoro-1,4-diazoniabicyclo[2,2,2]octane bis(tetrafluoroborate)] [9], montmorillonite K10 [10], $ZrOCl_2 \cdot 8H_2O$ [11], heteropolyacid supported MCM-41 [12], nanosized MCM-41- SO_3H [13], nano CuI [14], nano $SO_4^{2-}/xFe-Zr-O$ [15], silica bonded *N*-propyl sulfamic acid [16], $Zn[L\text{-proline}]_2$ [17],

[Bmim][HSO_4] [18], [Hmim]TFA [19], [TMPSA] HSO_4 [20], [DDPA][HSO_4] [21], [Et₃N- SO_3H Cl] [22], [Hbim]BF₄ [23], [Et₃NH]H₂PO₄ [24], [Et₃NH][HSO_4] [25], TMGT [26], *p*-dodecylbenzenesulphonic acid [27], trichloroisocyanuric acid [28], silica sulfuric acid [29], alumina-sulfuric acid [30], trimethylsilyl chloride [31], $NaHSO_4 \cdot SiO_2$ [32], ZnO-acetyl chloride [33], CAN [34], Amberlyst-15 [35], Dowex-50W [36], PMA- SiO_2 [37], PPA- SiO_2 [38], as catalysts and also in ionic liquid [bmim]BF₄ as solvent catalysed by $FeCl_3 \cdot 6H_2O$ [39], $InCl_3 \cdot 4H_2O$ [40], and $NaHSO_4$ [41], $Mg(BF_4)_2$ doped in [BMIm][BF₄] [42] and LiBr [43], $SmCl_3$ [44] etc. All the methods have their own advantages with some disadvantages such as low yields, requirement of organic solvents, use of expensive catalysts, tedious workup procedure, etc.

In recent years, ionic liquids are of considerable interest as they can serve as catalysts, green reaction media in organic synthesis and also due to their properties such as non-volatility, non-flammability, ionic conductivity [45-48]. Recently a variety of ionic liquids have been demonstrated as practical alternative to organic solvents as well as catalysts in organic reactions/transformations [49]. Recently, 1-(Benzenesulfonyl)-3-methyl-1*H*-imidazolium chlorides catalyzed electrophilic substitution reactions of indoles with various carbonyl compounds for an efficient one pot synthesis of

bis(indolyl)methanes have also been reported by our group^[50]. Also the use of microwave irradiation for carrying out organic reactions has been well established in recent years^[51-52] as it is an environmentally friendly method for the organic synthesis. herein, we report the catalytic application of 1-(3, 4-dichlorobenzenesulfonyl)-3-methyl-1*H*-imidazolium chlorides for the synthesis of 1,8-dioxo-octahydroxanthenes by the condensation of aromatic aldehydes with 5,5-dimethyl-1,3-cyclohexanedione using easily available, reusable ionic liquid as catalyst under microwave irradiation in aqueous media.

2. EXPERIMENTAL

All solvents and reagents were commercial and used without further purification. Melting points were uncorrected. Nuclear magnetic resonance spectra were obtained on a Bruker AMX spectrophotometer in CDCl₃ at 300 MHz. Chemical shifts were obtained in parts per million (δ) and were measured using tetramethyl silane (TMS) as reference. IR spectra were recorded on a Shimadzu FT-IR-8400S spectrophotometer using KBr pellets and were reported as wave numbers (ν cm⁻¹). The synthesis was carried out using modified household microwave oven (Frigidaire).

2.1. Preparation of [3,4-dcbismim][Cl]

1-Methyl-1*H*-imidazole (2.0 mmol) was taken in a dry round bottomed flask at 0°C. To this 3,4-dichlorobenzenesulfonyl chloride (2.0 mmol) was added slowly and the reaction mixture stirred for 3-4 min to obtain a light yellow precipitate. The product was washed with petroleum ether, dried and stored in cool place. The compound was obtained in 95-96 % yield.^[50]

2.2. General procedure for the microwave synthesis of 1,8-dioxo-octahydroxanthenes:

A mixture of aldehyde (5.0 mmol) and 5,5-dimethyl-1,3-cyclohexanedione (10 mmol), 1-(3, 4-dichlorobenzenesulfonyl)-3-methyl-1*H*-imidazolium chloride ([3,4-dcbismim][Cl]) (0.5 mmol) and 5 mL of water was taken in a closed teflon vessel. The mixture was irradiated with microwaves for a given time (Table 2). The reaction was monitored by TLC. After completion of the reaction, the product was filtered out from the water media and then re-crystallized from ethanol. Further, the filtrate (water) was vacuum evaporated to dryness to recover the catalyst [3,4-dcbismim][Cl] and the resulting catalyst was reused directly for the next run. The products were characterized by melting points, IR, NMR & Mass spectra and compared with those reported for the authentic samples.

2.3. Spectral Data for the few compounds

2.3.1. 9-phenyl-3,3,6,6-tetramethyl-1,2,3,4,6,7,8-octahydroxanthene-1,8-dione (3a)

White solid; m.p.=205-207°C. IR (KBr)(ν_{\max} /cm⁻¹): 2972,1670,1360,1196. ¹HNMR(CDCl₃,300MHz): δ =0.97(s,6H),1.08(s,6H),2.15-2.26(q, 4H), 2.48(s,4H), 4.72(s, 1H),7.07-7.14(m,1H),7.16-7.26(m,2H),7.36(m,2H). ¹³CNMR (CDCl₃100MHz): δ =27.2, 29.2, 31.9, 32.0, 41.3, 50.3, 114.6, 126.3, 128.0, 128.3, 144.0, 163.2, 195.0. MS (EI) *m/z*350 (M⁺).

2.3.2. 9-(4-chlorophenyl)-3,3,6,6-tetramethyl-1,2,3,4,6,7,8-octahydroxanthene-1,8-dione (3b)

White solid; m.p.=229-232°C. IR (KBr)(ν_{\max} /cm⁻¹): 2960,1660,1350,1190. ¹HNMR(CDCl₃,300MHz): δ =0.99(s,6H),1.08(s,6H),2.14-2.26(q, 4H), 2.46(s,4H), 4.70(s, 1H),7.16-7.42(m,4H). ¹³CNMR (CDCl₃100MHz): δ =27.2, 28.1, 30.8, 31.9, 32.1, 40.7, 50.6, 115.1, 128.1, 129.0, 129.6, 142.6, 162.4, 196.3. MS (EI) *m/z*385 (M+1).

2.3.3. 9-(4-hydroxyphenyl)-3,3,6,6-tetramethyl-1,2,3,4,6,7,8-octahydroxanthene-1,8-dione (3f)

White solid; m.p.=245-248°C. IR (KBr)(ν_{\max} /cm⁻¹): 3330,2954,1650,1360,1191. ¹HNMR(CDCl₃,300MHz): δ =0.99(s,6H),1.09(s,6H),2.14-2.20(q, 4H), 2.47(s,4H), 4.63(s, 1H),5.72(s,1H), 7.16-7.02(m,4H). ¹³CNMR (CDCl₃100MHz): δ =27.2, 29.1, 30.2, 31.4, 32.1, 40.6, 50.6, 115.1, 128.0, 129.0, 139.4, 140.6, 162.4, 195.3. MS (EI) *m/z*366 (M⁺).

3. RESULTS AND DISCUSSION


Initially, the catalyst [3,4-dcbismim][Cl] was prepared as previously explained. In order to optimize the green reaction conditions, we firstly investigated the greener solvents on the reaction rate as well as yields of the products (Table 1). In a typical experimental procedure, the reaction of 5,5-dimethyl-1,3-cyclohexanedione (1.0 mmol) and benzaldehyde (0.5 mmol) catalyzed by [3,4-dcbismim][Cl] (0.5 mmol) in different solvent media was investigated under microwaves. Water is found as an ideal solvent because it is inexpensive, non-flammable, abundant and environmentally benign.

To determine the optimum concentration of catalyst, we have investigated the model reaction of benzaldehyde (10.0 mmol) with 5, 5-dimethyl-1, 3-cyclohexanedione (20.0 mmol) carried out at 1, 5, 10, and 20 mol% of [3,4-dcbismim][Cl] in water under microwave irradiation. The products were obtained in about 90, 92, 94 and 94 % yield, respectively in 3min. This indicates that 10 mol% of [3,4-dcbismim][Cl] is sufficient for the best yield. Subsequently, all the

Table - 1: Effect of solvent in the preparation of 3,3,6,6-tetramethyl-9-phenyl-3,4,6,7-tetrahydro-2H-xanthene-1,8(5H,9H)-dione catalyzed by [3,4-dcbsmim][Cl].^a


Entry	Solvent	Catalyst (mol %)	Time (min)	Yield (%) ^b
1	Acetonitrile(3mL) + [Bmim][Cl] (7 mL)	10	4	93
2	Acetonitrile (10 mL)	10	4	94
3	[Bmim][Cl] (10 mL)	10	4	93
4	Water (10 mL)	10	3	94
5	Neat	10	4	90

^a Reaction conditions: benzaldehyde (0.5 mmol), 5,5-dimethyl-1,3-cyclohexanedione (1.0 mmol), Microwaves in different solvents; ^b Isolated yield.

**Scheme - 1: Synthesis of 1,8-dioxo-octahydroxanthenes catalyzed by [3,4-dcbsmim][Cl].****Table - 2: [3,4-dcbsmim][Cl] catalyzed microwave synthesis of 1,8-dioxo-octahydroxanthenes.^a**

Entry	Aldehyde	Product 3	Time (min)	Yield ^b (%)	Melting point (°C)
1	Benzaldehyde	3a	3	94	205-207
2	4-Chlorobenzaldehyde	3b	4	95	229-232
3	2-Chlorobenzaldehyde	3c	5	94	225-227
4	4-Bromobenzaldehyde	3d	5	90	240-242
5	4-Fluorobenzaldehyde	3e	5	91	226-228
6	4-Hydroxybenzaldehyde	3f	5	92	245-248
7	2-Hydroxybenzaldehyde	3g	4	89	202-206
8	4-Methoxybenzaldehyde	3h	5	90	240-242
9	3-Methoxybenzaldehyde	3i	5	95	178-180
10	4-Nitrobenzaldehyde	3j	3	92	224-226
11	3- Nitrobenzaldehyde	3k	5	93	170-172
12	4-Tolualdehyde	3l	4	90	208-209
13	2,4-Dichlorobenzaldehyde	3m	5	91	247-249
14	Cinnamaldehyde	3n	4	90	178-180
15	4-Hydroxy-3-methoxybenzaldehyde	3o	5	85	226-228

^a Reaction conditions: aromatic aldehyde (0.5 mmol), 5,5-dimethyl-1,3-cyclohexanedione (1.0 mmol), [3,4-dcbsmim][Cl] (0.5 mmol), Microwave irradiation, in aqueous medium; ^b Isolated yield.



Scheme - 2: Plausible mechanism for the synthesis of 1,8-dioxo-octahydroxanthenes catalyzed by [3,4-dcbsmim][Cl] in water.

Table - 3: Recycling of the catalyst.^a

Run	Time(min)	Yield ^b (%)
1	3	92
2	3	90
3	4	87
4	4	85

^a Reaction conditions: 4-nitro benzaldehyde (0.5 mmol), 5,5-dimethyl-1,3-cyclohexanedione (1.0 mmol), [3,4-dcbsmim][Cl] (0.5 mmol), Microwaves, in aqueous medium; ^b Isolated yield.

reactions (Table 2) were carried out under microwave in presence of 10 mol% of [3,4-dcbsmim][Cl] in aqueous conditions for the preparation of 1,8-dioxooctahydroxanthenes (3) (Scheme 1) and the expected product was obtained in good yield. Also at room temperature we obtained the first dehydration product of dimedone and aromatic aldehyde, that is, 2,2'-arylmethylenebis(3-hydroxy-5,5-dimethyl-2-cyclohexane-1-one (Intermediate A), as shown in scheme 2. The second dehydration step requires heating to remove water to afford the target 1,8-dioxooctahydroxanthenes (3).

In order to demonstrate the reusability of [3,4-dcbsmim][Cl] the experiment was performed using 4-nitrobenzaldehyde as a model substrate

under aqueous condition. The product was recrystallized from 95% ethanol to obtain the desired product in pure form. The water layer consisted of [3,4-dcbsmim][Cl] along with some residual reactants and byproducts, and vacuumed to get catalyst with impurities. The resulting catalyst was directly used in subsequent runs without further treatment (Table 3).

The ionic liquid [3,4-dcbsmim][Cl] can be recycled for at least 3-4 times without any loss in catalytic activity. The efficacy of [3,4-dcbsmim][Cl] in the synthesis of Xanthenes in comparison with the other ionic liquids as reported in the literature can be understood from the results shown in table 4.

Table - 4: Comparison of ionic liquids used as catalyst for the synthesis of 1,8-dioxo-octahydroxanthenes.^a

Entry	Ionic liquid	Conditions	Time (min)	Yield (%) ^b	Reference
1	[Bmim] HSO ₄	100 °C	25	93	18
2	TMGT/TFA	75 °C	20-30	92	26
3	[Et ₃ NH][H ₂ PO ₄]	100 °C	5	96	24
4	Mg(BF ₄) ₂ doped in [Bmim][BF ₄]	80 °C	30	97	42
5	[Et ₃ NH][HSO ₄]	100 °C	20	94	25
6	No Catalyst	Water, MW	7	80	-
7	[3,4-dcbsmim][Cl]	Water, MW	3	94	Present work

^a Reaction conditions: benzaldehyde (0.5 mmol), 5,5-dimethyl-1,3-cyclohexanedione (1.0 mmol) at different conditions; ^b Isolated yield.

4. CONCLUSION

In conclusion, 1-(3,4-dichlorobenzenesulfonyl)-3-methyl-1*H*-imidazolium chloride [3,4-dcbsmim][Cl] has proven to be an efficient and recyclable catalyst for the synthesis of xanthene derivatives under microwave irradiation. The method is simple, clean, and green. The recycled ionic liquid can be reused for 3-4 cycles without loss in catalytic activity.

Acknowledgements:

The authors are thankful to Bangalore University, Bangalore, for the financial support through YRB-BUIRF. We also thank NMRC, IISc., Bangalore for NMR spectral analysis and Department of Organic Chemistry IISc., Bangalore for mass spectral analysis.

5. REFERENCES

- Chibale K, Visser M, Schalkwyk D, Smith PJ, Saravanamuthu A and Fairlamb AH. Exploring the potential of xanthene derivatives as trypanothione reductase inhibitors and chloroquine potentiating agents. **Tetrahedron.**, 2003; 59: 2289-2296.
- Zarei A, Hajipour AR and Khazdooz L. The one-pot synthesis of 14-aryl or alkyl-14*H*-dibenzo[a,j]xanthenes catalyzed by P₂O₅/Al₂O₃ under microwave irradiation. **Dyes Pigments.**, 2010; 85: 133-138.
- Banerjee A and Mukherjee AK. Chemical aspects of santalin as a histological stain. **Stain Technology.**, 1981; 56: 83-85.
- Ellis GP. The chemistry of heterocyclic compounds, in chromenne, chromanes, and chromone. Edited by Wessberger A. and Taylor ECE. **John Wiley, NewYork**, 1977; 2: 13.
- Menchen SM, Benson SC, Lam JYL, Zhen W, Sun D, Rosenblum BB, Khan SH and Taing M. **US Patent US 6583168, 2003**; (Chem. Abstr. 139 (2003) 54287f).
- Licudine PJA, Kawate MK and Li QX. Method for the analysis of phloxine B, uranine and related xanthene dyes in soil using supercritical fluid extraction and high-performance liquid chromatography. **J. Agric. Food Chem.**, 1997; 45: 766-773.
- Knight CG and Stephens T. Xanthene-dye-labelled phosphatidylethanolamines as Probes of Interfacial pH. studies in phospholipid vesicles. **Biochem. J.**, 1989; 258: 683-689.
- Saint-Ruf G, Hieu HT and Poupelin JP. The effect of dibenzoxanthenes on the paralyzing action of zoxazolamine. **Naturwissenschaften.**, 1975; 62: 584-585.
- Heravi MRP. Selectfluor™ promoted synthesis of 9-aryl-1,8-dioxooctahydroxanthene derivatives under solvent-free conditions. **J. Iran. Chem. Soc.**, 2009; 6: 483-488.
- Dabiri M, Azimi SC and Bazgir A. One-pot synthesis of xanthene derivatives under solvent-free conditions. **Chem. Pap.**, 2008; 62: 522-526.
- Lü HY, Li JJ and Zhang ZH. ZrOCl₂ · 8H₂O: a highly efficient catalyst for the synthesis of 1,8-dioxo-octahydroxanthene derivatives under solvent-free conditions. **Appl. Organomet. Chem.**, 2009; 23:165-169.
- Karthikeyana G and Pandurangana A. Heteropolyacid (H₃PW₁₂O₄₀) supported MCM-41: An efficient solid acid catalyst for the green synthesis of xanthenedione derivatives. **J. Mol. Catal. A: Chem.**, 2009; 311: 36-45.

13. Rostamizadeh S, Amani AM, Mahdavinia GH, Amiri G and Sepehrian H. Ultrasound promoted rapid and green synthesis of 1,8-dioxo-octahydroxanthenes derivatives using nanosized MCM-41-SO(3)H as a nanoreactor, nanocatalyst in aqueous media. **Ultrason. Sonochem.**, 2010; 17: 306-309.
14. Albadi J, Keshavarz M, Abedini M and Khoshakhlagh M. Copper iodide nanoparticles on Poly (4-vinylpyridine): a new and efficient catalyst for the synthesis of 1,8-dioxooctahydroxanthenes under solvent-free conditions. **J. Chem. Sci.**, 2013; 125: 295-298.
15. Samantaray S, Kar P, Hota G and Mishra BG. Sulfate grafted iron stabilized zirconia nanoparticles as efficient heterogenous catalysts for solvent-free synthesis of xanthenediones under microwave irradiation. **Ind. Eng. Chem. Res.**, 2013; 52: 5862-5870.
16. Rashedian F, Saberi D and Niknam K. Silica-bonded *N*-propyl sulfamic acid: a recyclable catalyst for the synthesis of 1, 8-dioxo-decahydroacridines, 1,8-dioxo-octahydroxanthenes and quinoxalines. **J. Chin. Chem. Soc.**, 2010; 57: 998-1006.
17. Kidwai M and Jain A. Zn [(L)Proline] 2: an eligible candidate for the synthesis of xanthenediones in water. **Appl. Organomet. Chem.**, 2012; 26: 528-535.
18. Niknam K and Damya M. 1-Butyl-3-methylimidazolium hydrogen sulfate [Bmim]HSO₄: an efficient reusable acidic ionic liquid for the synthesis of 1,8-dioxo-octahydroxanthenes. **J. Chin. Chem. Soc.**, 2009; 56: 659-665.
19. Dabirir M, Baghbanzadeh M and Arzroomchilar E. 1-Methylimidazolium trifluoroacetate ([Hmim]TFA): An efficient reusable acidic ionic liquid for the synthesis of 1,8-dioxo-octahydroxanthenes and 1,8-dioxo-decahydroacridines **Catal. Commun.**, 2008; 9: 939-942.
20. Fang D, Gong K and Liu ZL. Synthesis of 1,8-dioxo-octahydroxanthenes catalyzed by acidic ionic liquids in aqueous media. **Catal. Lett.**, 2008; 127: 291-295.
21. Fang D, Yang J and Liu Z. Eco-friendly synthesis of 1,8-dioxo-octahydroxanthenes catalyzed by ionic liquid in aqueous media. **J. Heterocycl. Chem.**, 2011; 48: 468-472.
22. Zare A, Moosavi-Zare AR, Merajoddin M, Zolfigol MA, Hekmat-Zadeh T, Hasaninejad A and Khazaei A. Ionic Liquid Triethylamine-bonded sulfonic acid {[Et₃N-SO₃H]Cl} as a novel, highly efficient and homogeneous catalyst for the synthesis of β-acetamido ketones, 1,8-dioxo-octahydroxanthenes and 14-aryl-14H-dibenzo[a,j]xanthenes. **J. Mol. Liq.**, 2012; 167: 69-77.
23. Venkatesan K, Pujari SS, Lahoti RJ and Srinivasan KV. An efficient synthesis of 1,8-dioxooctahydroxanthene derivatives promoted by a room temperature ionic liquid at ambient conditions under ultrasound irradiation. **Ultrason. Sonochem.**, 2008; 15: 548-553.
24. Kalantari M. Synthesis of 1, 8-dioxo-octahydroxanthenes and bis(indolyl)methanes catalyzed by [Et₃NH][H₂PO₄] as a cheap and mild acidic ionic liquid. **Arabian J. Chem.**, 2012; 5:319-323.
25. Zhou Z and Deng X. [Et₃NH][HSO₄] catalyzed efficient and green synthesis of 1,8-dioxooctahydroxanthenes. **J. Mol. Catal. A: Chem.**, 2013; 367: 99-102.
26. Rahmati A. A rapid and efficient method for the synthesis of 14H-dibenzo[α,j] xanthenes, aryl-5H-dibenzo [b.i]xanthene-5,7,12,14-(13H)-tetraone and 1,8-dioxo-octahydroxanthenes by acidic ionic liquid. **Chin. Chem. Lett.**, 2010; 21: 761-764.
27. Jin T, Zhang J, Xiao J, Wang A and Li T. Clean synthesis of 1,8-dioxooctahydroxanthene derivatives catalyzed by *p*-dodecylbenzenesulfonic acid in aqueous media. **Synlett.**, 2004; 0866-0870.
28. Bigdeli MA, Nemati F, Mahdavinia GH and Doostmohammadi H. A series of 1,8-dioxooctahydroxanthenes are prepared using trichloroisocyanuric acid. **Chin. Chem. Lett.**, 2009; 20: 1275-1278.
29. Seyyedhamzeh M, Mirzaei P and Bazgir A. Solvent-free synthesis of aryl-14H-dibenzo [a,j]xanthenes and 1,8-dioxooctahydroxanthenes using silica sulfuric acid as catalyst. **Dyes Pigments.**, 2008; 76: 836-839.
30. Pramanik A and Bhar S. Alumina-sulfuric acid catalyzed eco-friendly synthesis of xanthenediones. **Catal. Commun.**, 2012; 20:17-24.
31. Kantevari S, Bantu R and Nagarapu L. TMSCl mediated highly efficient one-pot synthesis of octahydroquinazolinone and 1, 8-dioxooctahydroxanthene derivatives. **ARKIVOC.**, 2006; part XVI: 136-148.
32. Das B, Thirupathi P, Reddy KR, Ravikanth B and Nagarapu L. An efficient synthesis of 1,8-dioxooctahydroxanthenes using heterogeneous catalysts. **Catal. Commun.**, 2007; 8: 535-538.

33. Maghsoodlou MT, Habibi-Khorassani SM, Shahkarami Z, Maleki N and Rostamizadeh M. An efficient synthesis of 2, 2'-arylmethylene bis(3-hydroxy-5,5-dimethyl-2-cyclohexene-1-one) and 1,8-dioxooctahydroxanthenes using ZnO and ZnO-acetyl chloride. **Chin. Chem. Lett.**, 2010; 21: 686-689.
34. Sudha S and Pasha MA. Ultrasound assisted synthesis of tetrahydrobenzo[c]xanthene-11-ones using CAN as catalyst. **Ultrason. Sonochem.**, 2012; 19: 994-998.
35. Das B, Thirupathi P, Mahender I, Reddy VS and Rao YK. Amberlyst-15: an efficient reusable heterogeneous catalyst for the synthesis of 1,8-dioxooctahydroxanthenes and 1,8-dioxodecahydroacridines. **J. Mol. Catal. A: Chem.**, 2006; 247: 233-239.
36. Shakibaei GI, Mirzaei P, Bazgir A. Dowex-50W promoted synthesis of 14-aryl-14H-dibenzo[a,j]xanthene and 1,8-dioxo-octahydroxanthene derivatives under solvent-free conditions. **Appl. Catal. A: Gen.** 2007; 325:188-192.
37. Srihari P, Mandal SS, Reddy JSS, Rao RS, Yadav JS. Synthesis of 1,8-dioxo-octahydroxanthenes utilizing PMA-SiO₂ as an efficient reusable catalyst. **Chin. Chem. Lett.**, 2008; 19: 771-774.
38. Kantevari S, Bantu R and Nagarapu L. HClO₄-SiO₂ and PPA-SiO₂ catalyzed efficient one-pot Knoevenagel condensation, Michael addition and cyclodehydration of dimedone and aldehydes in acetonitrile, aqueous and solvent free conditions: scope and Limitations. **J. Mol. Catal. A: Chem.**, 2007; 269: 53-57.
39. Fan X, Li Y, Zhang X, Hu X and Wang J. FeCl₃ • 6H₂O catalyzed condensation of aromatic aldehydes with 5, 5-dimethyl-1, 3-cyclohexanedione in ionic liquids. **Chin. J. Org. Chem.**, 2005; 25: 1482-1486.
40. Fan X, Hu X, Zhang X and Wang J. InCl₃.4H₂O-promoted green preparation of xanthenedione derivatives in ionic liquids. **Can. J. Chem.**, 2005; 83: 16-20.
41. Ma J, Li J, Tang R, Zhou X, Wu Q and Wang C. Condensation of aromatic aldehydes with 1,3-cyclohexanedione catalyzed by NaHSO₄ in ionic liquids. **Chin. J. Org. Chem.**, 2007; 27: 640-642.
42. Rad-Moghadam K and Azimi SC. Mg(BF₄)₂ Doped in [BMIm][BF₄]: a homogeneous ionic liquid-catalyst for efficient synthesis of 1,8-dioxo-octahydroxanthenes, decahydroacridines and 14-aryl-14H-dibenzo[a,j]xanthenes. **J. Mol. Catal. A: Chem.**, 2012; 363-364: 465-469.
43. Kumar D and Sandhu JS. Efficient, solvent-free, microwave-enhanced condensation of 5,5-dimethyl-1,3-cyclohexanedione with aldehydes and imines using LiBr as inexpensive, mild catalyst. **Synth. Commun.**, 2010; 40: 510-517.
44. Ilangovan A, Malayappasamy S, Muralidharan S and Maruthamuthu S. A highly efficient green synthesis of 1, 8-dioxo-octahydroxanthenes. **Chemistry Central Journal.**, 2011; 5: 81.
45. MacFarlane DR and Seddon KR. Ionic Liquids-progress on the fundamental issues. **Aust. J. Chem.**, 2007; 60: 3-5.
46. Clare B, Sirwardana A and Macfarlane DR. Synthesis, purification and characterization of ionic liquids. Springer-Verlag, Berlin-Heidelberg. 2009; 290: 1-40.
47. Bonhôte P, Dias A-P, Papageorgiou N, Kalyanasundaram K and Gratzel M. Hydrophobic, highly conductive ambient-temperature Molten Salts. **Inorg. Chem.**, 1996; 35: 1168-1178.
48. Huddleston JG, Visser AE, Reichert WM, Willauer HD, Broker GA and Rogers RD. Characterization and comparison of hydrophilic and hydrophobic room temperature ionic liquids incorporating the imidazolium cation. **Green Chem.**, 2001; 3:156-164.
49. Welton T. Room-temperature ionic liquids. Solvents for synthesis and catalysis. **Chem. Rev.**, 1999; 99: 2071-2083.
50. Krishnamurthy G and Jagannath KV. Efficient green synthesis of bis(3-indolyl) phenylmethanes catalyzed by 1-(Benzenesulfonyl)-3-methyl-1H-imidazolium chlorides. **Letters in Organic Chemistry.**, 2013; 10: 744-751.
51. Kappe CO, Stadler A and Dallinger D. Microwaves in organic and medicinal chemistry. **Wiley-vch, Weinheim.** 2012; 52.
52. Krishnamurthy G and Jagannath KV. Microwave-assisted Silica-promoted Solvent-free Synthesis of Triazoloquinazolinone and Benzimidazoquinazolinones. **J. Chem. Sci.**, 2013; 125: 807-811.